

Just 25 minutes (+5 minutes for questions) to save the world!



THE BIRCHALL CENTRE

The
Birchall
CENTRE



Innovations in Inorganic and Materials Chemistry

Aluminium and Alzheimer's Disease

Christopher Exley PhD FRSB


The Birchall Centre, Lennard-Jones Laboratories, Keele University, Keele,
Staffordshire, ST5 5BG, UK.

c.exley@keele.ac.uk

<http://www.keele.ac.uk/aluminium/>









Achieving Optimal Health, Washington
2016




THE BIOGEOCHEMICAL CYCLE OF ALUMINIUM

Exley C (2003) *J. Inorg. Biochem.* 97, 1-7.





researching the cure




THE ALUMINIUM AGE?


<http://www.youtube.com/watch?v=FrYxIVb3qPs>

AUTISM
Awareness



Unmasking Osteomalacia






Achieving Optimal Health, Washington

2016

Alzheimer's Disease







Figure 1. The sequence of major pathogenic events leading to AD proposed by the amyloid cascade hypothesis.

The curved blue arrow indicates that Aβ oligomers may directly injure the synapses and neurites of brain neurons, in addition to activating microglia and astrocytes.



Why industry propaganda and political interference cannot disguise the inevitable role played by human exposure to aluminum in neurodegenerative diseases, including Alzheimer's disease

Christopher Exley *

The Birchall Centre, Lennard-Jones Laboratories, Keele University, Stoke-on-Trent, UK

Edited by:

Christopher Ariel Shaw, University of British Columbia, Canada

Reviewed by:

Nelson Silva Filho, Universidade Estadual Paulista, Brazil
Lucija Tomljenovic, University of British Columbia, Canada

***Correspondence:**

Christopher Exley, The Birchall Centre, Lennard-Jones Laboratories, Keele University, Staffordshire ST5 5BG, UK
e-mail: c.exley@keele.ac.uk

In the aluminum age, it is clearly unpalatable for aluminum, the globe's most successful metal, to be implicated in human disease. It is unpalatable because for approximately 100 years human beings have reaped the rewards of the most abundant metal of the Earth's crust without seriously considering the potential consequences for human health. The aluminum industry is a pillar of the developed and developing world and irrespective of the tyranny of human exposure to aluminum it cannot be challenged without significant consequences for businesses, economies, and governments. However, no matter how deep the dependency or unthinkable the withdrawal, science continues to document, if not too slowly, a burgeoning body burden of aluminum in human beings. Herein, I will make the case that it is inevitable both today and in the future that an individual's exposure to aluminum is impacting upon their health and is already contributing to, if not causing, chronic diseases such as Alzheimer's disease. This is the logical, if uncomfortable, consequence of living in the aluminum age.

Keywords: aluminium, Alzheimer's disease, human exposure, neurodegenerative disease, body burden

Cite this: *Metallomics*, 2012, **4**, 56–65

www.rsc.org/metallomics

PAPER

Aluminium, iron and copper in human brain tissues donated to the medical research council's cognitive function and ageing study

Emily House,^a Margaret Esiri,^b Gill Forster,^c Paul G Ince^c and Christopher Exley*^a

Sixty Human Brains

>700 tissue digests (250 mg wet weight)

174 Method Blanks; Al = 54 ng Al/vessel (Mean + 1.654SD)

House et al. (2012) *Metallomics* 4, 56.

Achieving Optimal Health, Washington

2016

Aluminium Content of Human Brain




Fig. 3 Percentage frequency (bars) and cumulative frequency (line and marker) distributions of (a) Fe, (b) Cu and (c) Al concentrations in ($n = 719, 720$ and 713 respectively) brain tissues after subtraction of contamination.

The median Al content of tissues from all **60 brains** ($n=713$) is $1 \mu\text{g/g}$ dry wt.

In 52 out of 60 individuals at least one tissue sample exceeded $2 \mu\text{g Al/g}$ dry wt.

In 41 out of 60 individuals at least one tissue sample exceeded $3.5 \mu\text{g Al/g}$ dry wt.

Approximately 70% of individuals aged 70 – 103 years had at least one tissue Al content which should be considered as pathological.

Accidental Exposure to Aluminium

Camelford, Cornwall, United Kingdom, 1988

SHORT REPORT


Severe cerebral congophilic angiopathy coincident with increased brain aluminium in a resident of Camelford, Cornwall, UK

C Exley, M M Esiri



See Editorial Commentary, p 811

J Neurol Neurosurg Psychiatry 2006;77:877-879. doi: 10.1136/jnnp.2005.086553



Frontal Cortex, n=5.

1. 23.00* $\mu\text{g/g}$ dry wt.
2. 3.24
3. 11.01
4. 4.33
5. 5.71

*Noted during measurement as a heavily vascularised piece of tissue.

Exley & Esiri (2006) *J Neurol Neurosurg Psychiatr* 77, 877.

Achieving Optimal Health, Washington

2016

Occupational Exposure to Aluminium

Exley and Vickers *Journal of Medical Case Reports* 2014, **8**:41
<http://www.jmedicalcasereports.com/content/8/1/41>



CASE REPORT

Open Access

Elevated brain aluminium and early onset Alzheimer's disease in an individual occupationally exposed to aluminium: a case report

Mean Al content of frontal lobe tissue (n=46) is 2.98 (2.73) $\mu\text{gAl/g}$ dry wt.

Range is 0.00 (less than the method blank) to 12.97 $\mu\text{gAl/g}$ dry wt.

More than 30% of tissue samples had an Al content considered as pathological, greater than 3.50 $\mu\text{gAl/g}$ dry wt.

Another case of early onset ‘sporadic’ Alzheimer’s disease from Camelford

Table 1

The aluminum content of each tissue replicate from each brain region ($\mu\text{g/g}$ dry wt.) with pathologically-concerning values highlighted in **bold** and pathologically-significant values highlighted in **bold italics**. Mean and SD are also given for each region for between 9 and 21 replicates

Replicate	Temporal	Hippocampus	Occipital	Frontal	Parietal
1	0.81	0.41	4.45	1.31	1.21
2	0.12	0.01	2.68	0.40	2.05
3	3.51	2.73	0.09	0.77	1.84
4	0.66	0.76	0.19	1.01	0.84
5	0.01	0.41	1.19	0.27	0.04
6	0.74	0.01	0.57	1.89	0.36
7	1.18	0.11	0.24	0.48	2.78
8	0.24	0.01	2.82	0.01	5.58
9	1.71	0.01	0.39	0.02	0.41
10		0.01	1.03	0.32	0.98
11			0.27	0.27	0.01
12			0.72	0.26	0.01
13			2.79	0.28	1.23
14			4.00	0.14	0.93
15			0.58	0.39	0.29
16			1.56	0.20	0.34
17			2.16	0.69	2.62
18			3.73	0.43	0.31
19			2.16	0.20	3.91
20			1.23	3.38	1.33
21			3.86	1.03	0.79
Mean	1.00	0.45	1.75	0.66	1.33 (1.42)
(SD)	(1.08)	(0.84)	(1.43)	(0.78)	(1.42)

What about; Familial Alzheimer's Disease?

Summary indicating the lobe with the highest content of AI (Mean (SD) n=3)

(Pathologically Concerning/Pathologically Significant / 12 tissues)

Donor ID	Lobe	[AI] µg/g dry wt.
A1 0/0 /12	Frontal	0.44(0.31)
A2 0/4 /12	Occipital	9.99(1.61)
A3 6/3 /12	Occipital	4.29(3.21)
A4 3/2 /12	Parietal	3.30(4.43)
A5 1/2 /12	Frontal	7.05(6.10)
A6 1/3 /12	Occipital	9.57(14.08)
A7 2/0 /12	Temporal	1.81(0.78)
A8 3/6 /12	Frontal	14.41(18.44)
A9 4/2 /12	Frontal	3.81(4.11)
A10 5/1 /12	Occipital	2.97(1.01)
A11 2/2/12	Occipital	9.31(12.71)
A12 2/3 /12	Temporal	2.91(2.89)

Mirza et al (submitted).

From the 'Discussion'

“The aluminium content of brain tissue donated by individuals with a diagnosis of familial AD was, overall, extremely high. We measured some of the highest values recorded for human brain tissues, for example to highlight just a few, 11.54 $\mu\text{g/g}$ in the occipital lobe of donor A2, 13.41 $\mu\text{g/g}$ in the frontal lobe of A5, 25.80 $\mu\text{g/g}$ in the occipital lobe of A6, 35.65 $\mu\text{g/g}$ in the frontal lobe of A8 and 23.93 $\mu\text{g/g}$ in the occipital lobe of A11 (Table 1)”

Another case of early onset sporadic Alzheimer's disease in a Camelford resident!

Journal of Alzheimer's Disease xx (20xx) x–xx
DOI 10.3233/JAD-160648
IOS Press

1

The Identification of Aluminum in Human Brain Tissue Using Lumogallion and Fluorescence Microscopy


Ambreen Mirza^a, Andrew King^{b,c}, Claire Troakes^c and Christopher Exley^{a,*}

^a*The Birchall Centre, Lennard-Jones Laboratories, Keele University, Staffordshire, UK*

^b*Department of Clinical Neuropathology, King's College Hospital, London, UK*


^c*MRC London Neurodegenerative Diseases Brain Bank, Institute of Psychiatry, Psychology and Neuroscience, King's College, London, UK*

Mirza et al. (2016) J Alzh Dis (In the press)



Frontal Cortex


Mirza et al (submitted).



Parietal Cortex


Mirza et al (submitted).

Achieving Optimal Health, Washington
2016



Frontal (A-C) and Temporal D-F) Cortex

Mirza et al (submitted).
Achieving Optimal Health, Washington
2016



Occipital Cortex

Mirza et al (submitted).

Achieving Optimal Health, Washington
2016

“Aluminium is neurotoxic and the concentrations of aluminium found in these familial AD brains are unlikely to be benign and indeed are highly likely to have contributed to both the onset and the aggressive nature of any ongoing AD. These data lend support to the recent conclusion that brain aluminium will contribute towards all forms of AD under certain conditions.”

Mirza et al. submitted

Vancouver, British Columbia



12th Keele Meeting on Aluminium Living in the Aluminium Age

Saturday, March 4 to Wednesday, March 8, 2017

Pinnacle Hotel, Vancouver



Keele
University



University of British Columbia
Neural Dynamics
Research Group



CHILDREN'S MEDICAL SAFETY
RESEARCH INSTITUTE



Information: www.keele.ac.uk; c.exley@keele.ac.uk;
cashawlab@gmail.com; suarez@mail.ubc.ca